

Tim Wang, M.D. Orthopaedic Surgery and Sports Medicine 10710 N. Torrey Pines Road 2205 Vista Way, Suite 210

La Jolla, CA 92037 Phone: 858-554-7980 Fax: 858-404-1813 2205 Vista Way, Suite 210 Oceanside, CA 92054 Phone: 760-704-5750 Fax: 858-404-1813

www.timwangmd.com

AC Joint Reconstruction

<u>Postop</u>	<u>Goals</u>	<u>Precautions</u>	<u>Exercises</u>
Weeks 0-4 PT 1x/week HEP daily	Edema and pain control Protect surgical repair Sling immobilization Keep elbow supported!	Sling at all times including sleep Remove for hygiene & exercises	Sidelying scapular stabilization Elbow, wrist, hand ROM Gripping exercises
Weeks 4-8 PT 1-2x/week HEP daily	Protect surgical repair Restore ROM	Sling except shower & exercises No ER >45° until 6 weeks No FF >120° until 6 weeks	Sidelying scapular stabilization Supine deltoid, biceps, triceps, rotator cuff isometrics (gravity eliminated) Pendulums/codmans Supine PROM: FF 120°, ER 45°
Weeks 8-16 PT 2-3x/week HEP daily	Full ROM Normalize scapulohumeral rhythm throughout ROM Restore strength 5/5	DC sling at 8 weeks Avoid painful ADL's Avoid rotator cuff inflammation Avoid excessive passive stretching OK to cycle/run at 12 weeks	AROM as tolerated in PRONE position AA/PROM no limits Continue scapular stabilization Begin resistive strengthening for scapula, biceps, triceps, and rotator cuff (supine → vertical position at 12 weeks) Endurance: Begin UBE at 12 weeks
Weeks 16-24 PT 1-2x/week HEP daily	Full ROM and strength Improve endurance Prevent re-injury	Avoid painful activities No contact/racket/throwing sports Return to sport (MD directed)	Advance eccentric training Initiate plyometrics Advance endurance training Sport specific activities Throwing/racquet program ~5 months Contact sports 6+ months